



LECTURE

Ataxin-3 phosphorylation decreases neuronal defects in spinocerebellar ataxia type 3 models

Carlos Andrade Matos^{1,2}, Clévio Nóbrega¹, Susana R. Louros¹, Bruno Almeida³, Elisabete Ferreiro^{1,4}, Jorge Valero^{1,5}, Luís Pereira de Almeida^{1,6}, Sandra Macedo-Ribeiro³ and Ana Luísa Carvalho^{1,2}

Special Issue on Controversies in Neurology. From the 10th World Congress on Controversies in Neurology (CONy), Lisbon, Portugal. 17–20 March 2016.

Abstract

Different neurodegenerative diseases are caused by aberrant elongation of repeated glutamine sequences normally found in particular human proteins. Although the proteins involved are ubiquitously distributed in human tissues, toxicity targets only defined neuronal populations. Changes caused by an expanded polyglutamine protein are possibly influenced by endogenous cellular mechanisms, which may be harnessed in order to produce neuroprotection. Here, we show that ataxin-3, the protein involved in Spinocerebellar ataxia type 3, also known as Machado-Joseph disease, causes dendritic

and synapse loss in cultured neurons when expanded. We report that S12 of ataxin-3 is phosphorylated in neurons and that mutating this residue so as to mimic a constitutive phosphorylated state counters the neuromorphologic defects observed. In rats stereotaxically injected with expanded ataxin-3- encoding lentiviral vectors, mutation of serine 12 reduces aggregation, neuronal and synapse loss. Our results suggest that S12 plays a role in the pathogenic pathways mediated by polyglutamine-expanded ataxin-3 and that phosphorylation of this residue protects against toxicity.

¹CNC – Center for Neuroscience and Cell Biology, University of Coimbra, 3004-504 Coimbra, Portugal

²Department of Life Sciences, Faculty of Sciences and Technology University of Coimbra, 3004-517 Coimbra, Portugal

³IBMC-Instituto de Biologia Molecular e Celular and i3S – Instituto de Investigação e Inovação em Saúde, University of Porto, 4200-135 Porto, Portugal

⁴Institute for Interdisciplinary Research, University of Coimbra, Coimbra, Portugal

⁵Ikerbasque Basque Foundation for Science and Achucarro Basque Center for Neuroscience, Bizkaia Science and Technology Park, E-48170 Zamudio, Spain

⁶Faculty of Pharmacy, University of Coimbra, 3000-548 Coimbra, Portugal

Citation: Matos et al. Ataxin-3 phosphorylation decreases neuronal defects in spinocerebellar ataxia type 3 models. *International Journal of Clinical Neurosciences and Mental Health* 2016; 3(Suppl. 1):L26

Published: 16 March 2016



Open Access Publication Available at <http://ijcnmh.arc-publishing.org>

© 2016 Matos et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

