



LECTURE

Mechanisms of disease: proteolysis of ataxin-3

Jonasz Jeremiasz Weber^{1,2}

Special Issue on Controversies in Neurology. From the 10th World Congress on Controversies in Neurology (CONy), Lisbon, Portugal. 17–20 March 2016.

Abstract

Proteolytic cleavage of disease-causing proteins is a widely discussed mechanism in neurodegenerative disorders, which gives rise to toxic fragments and thus may amplify the formation of protein aggregates. This molecular model - known as the toxic fragment hypothesis - was also proposed to play a crucial role in Machado Joseph Disease (MJD) and several studies have reported on the proteolysis of polyglutamine-expanded ataxin-3, the causative protein in MJD. Hitherto two classes of enzymes, caspases and calpains, were

linked with the proteolytic processing of ataxin-3, triggering neurotoxicity and cell death. Inhibition of protease activity and mutation of caspase or calpain cleavage sites within mutant ataxin-3 was shown to reduce toxicity and neuronal death in cell and animal models of MJD. Our better understanding of proteolytic events in MJD has opened new possibilities to identify therapeutic targets as a treatment for this fatal disorder.

¹Department of Medical Genetics and Applied Genomics, University of Tübingen, Tübingen, Germany

²Rare Disease Center, University of Tübingen, Tübingen, Germany

Citation: Weber, JJ. Mechanisms of disease: proteolysis of ataxin-3. International Journal of Clinical Neurosciences and Mental Health 2016; 3(Suppl. 1):L34

Published: 16 March 2016



Open Access Publication Available at <http://ijcnmh.arc-publishing.org>

© 2016 Weber, JJ. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

